

INFORMATION PAPER

DHA-IHD
15 November 2024

SUBJECT: Meningococcal Disease and Meningococcal Vaccines

1. Purpose. To describe meningococcal disease and the vaccines to prevent it.
2. Facts.
 - a. Microbiology. *Neisseria meningitidis*, or meningococcus, is an aerobic, gram-negative diplococcus, closely related to *N. gonorrhoea* and to several nonpathogenic *Neisseria* species, such as *N. lactamica*. *N. meningitidis* has both an inner and outer membrane, separated by a cell wall. The outer membrane contains several protein structures that enable the bacteria to interact with the host cells as well as perform other functions. The outer membrane is surrounded by a polysaccharide capsule that protects the organism from phagocytosis and complement-mediated lysis. There are thirteen distinct serogroups, which are based on the characteristics of the polysaccharide capsule. However, only 6 serogroups of *N. meningitidis* (A, B, C, W, X and Y) cause the majority of disease worldwide, but serotypes B, C, and Y cause most of the illness in the United States.
 - b. Disease. *N. meningitidis* colonizes mucosal surfaces of the nasopharynx and is transmitted through direct contact with respiratory droplet secretions from infected individuals and asymptomatic carriers. Humans are the only host. Meningococcal disease is a serious health threat, causing meningitis (inflammation of membranes around the brain and spinal cord) or blood infections (meningococemia). Bacterial meningitis can be extremely severe resulting in brain damage, hearing loss, or learning disability. For bacterial meningitis, it is important to know which type of bacteria is causing the meningitis because antibiotics can prevent some types from spreading and infecting other people. Common symptoms in individuals aged 2 years and older, which develop over several hours, or can take 1 to 2 days, include high fever, headache, and stiff neck. Other symptoms include nausea, vomiting, confusion, sleepiness, and discomfort looking into bright lights. In newborns and small infants, the classic symptoms may be difficult to detect. The infant may appear slow, inactive, and/or irritable, experience vomiting, or loss of appetite. As the disease progresses, individuals of any age may develop seizures. Meningococcal disease can be disfiguring or disabling (i.e., limb amputations, hearing loss, brain damage) in up to 20% of those who recover.

- c. Epidemiology. Meningococcal disease occurs worldwide in both endemic and epidemic form. Despite the use of effective antibiotics, meningococcal disease still results in death for 10% to 14% of those who become ill. Of note, outbreak-associated cases are associated with a higher case-fatality rate than sporadic cases (21% vs. 11%). Prior to the availability of the monovalent serotype A meningococcal conjugate vaccine in 2010 (MenAfriVac), 90% of cases in Africa (“meningitis belt”) were due to serotype A. More recently, cases in this region have been primary due to serogroups C and W. In the United States serotype B accounts for about 60% of meningococcal disease in children and young adults under the age of 25; and serotypes C, Y, and W-135 cause about 65% of illness in persons aged 25 years and older. Serious (also called invasive) meningococcal disease occurs most often in infants younger than 1 year of age and surges a second time in adolescence. High-risk groups include college freshmen and military trainees living in dormitories (likely due to crowded living conditions), people with immune deficiencies, travelers to areas where the disease is endemic (sub-Saharan Africa), and people who do not have a spleen or whose spleen is not functioning (as in sickle-cell anemia).
- d. Vaccines. There are three types of meningococcal vaccines available in the United States:
- (1) Serogroup ACWY Meningococcal conjugate vaccines:
- Menveo®: 2 vial presentation: 2 months-55 years
1 vial presentation: 10 years-55 years
 - MenQuadfi®: 2 years and older
 - a. Routinely, all healthy 11–12-year-olds should be vaccinated with a quadrivalent (protects against serogroups A, C, W, and Y) meningococcal conjugate vaccine. A booster dose is recommended at age 16 years. For adolescents who receive the first dose at age 13 through 15 years, a booster dose should be administered, preferably at age 16 through 18 years, before the period of increased risk. Adolescents who receive their first dose of quadrivalent meningococcal conjugate vaccine at or after age 16 years do not need a booster dose.
 - b. When quadrivalent meningococcal conjugate vaccine was first recommended for adolescents in 2005, the

expectation was that protection would last for 10 years; however, currently available data suggest it wanes in most adolescents within 5 years. Based on that information, a single dose at the recommended age of 11 or 12 years may not offer protection through the adolescent years at which risk for meningococcal infection is highest (16 through 23 years of age).

- c. For patients who are about to start college and got their first dose of quadrivalent meningococcal conjugate vaccine more than 5 years ago, it is recommended that these patients receive a booster dose of quadrivalent meningococcal conjugate vaccine.
- d. For patients younger than 16 years who require fewer healthcare visits, clinical judgment is recommended when determining when to provide the booster dose. The minimum interval between doses is 8 weeks.
- e. Adolescents who receive their first dose of meningococcal conjugate vaccine at or after age 16 years do not need a booster dose.
- f. Persons 2 months and older who are at increased risk for meningococcal disease should be vaccinated with the age appropriate MenACWY vaccine. Recommended vaccine, doses, and intervals vary according to age and target group.

Increased risk includes:

- i. Persons with complement component deficiency (e.g., C5-C9, properdin, factor H, factor D, or taking a complement inhibitor (e.g., eculizumab (Soliris®), ravulizumab (Ultomiris®))
- ii. Persons with functional or anatomic asplenia.
- iii. Microbiologist routinely exposed to *Neisseria meningitidis* (the causal pathogen).

- iv. Persons traveling or residing in countries in which the disease is common.
 - v. Persons identified to be at increased risk because of a serogroup A, C, W or Y meningococcal disease outbreak.
 - vi. First-year college student living in a residence hall.
 - vii. Military recruits.
- g. Infants 2-23 months of age with asplenia or HIV may receive Menveo without potential interference in the PCV response. MenQuadfi® (≥ 2years) was not studied in children but has no PCV impact in adolescents.
- h. Menveo® and MenQuadfi® (≥2 years) may be administered to a child at increased risk for meningococcal disease any time before or after DTaP.
- i. Although off-label, the ACIP recommends that individuals 56 years or older who require meningococcal vaccination because they are at increased risk for meningococcal disease (as listed above), receive either MenQuadfi® or Menveo®.

***Menactra® will no longer available to order in the United States after 2022 but may be used until the expiration date. Menomune® has not been available since September 2017. MenHibrix®, another meningococcal conjugate vaccine at one time used for high-risk individuals 2–15 months of age, was discontinued February 2017.*

(2) Serogroup B meningococcal vaccines (Bexsero® and Trumenba®):

- a. Routine vaccination recommended for people 10 years and older identified as being at increased risk. Increased risk includes:
 - i. Persons identified as at increased risk because of a serogroup B meningococcal disease outbreak.
 - ii. Microbiologists routinely exposed to *Neisseria meningitides* isolates.
 - iii. Persons with persistent complement component deficiencies (e.g., C5-C9, properdin, factor H, factor D), or are taking eculizumab (Soliris®), ravulizumab (Ultomiris®), or other complement inhibitor medications.
 - iv. Persons with anatomic or functional asplenia.
- b. Adolescents and young adults 16 through 23 years of age not at increased risk may be vaccinated with either MenB vaccine to provide short-term protection against most strains of serogroup B meningococcal disease based on shared clinical decision making.
 - i. MenB vaccines may be prescribed based on shared decision making for healthy first-year college students living in residence halls, military recruits, or other adolescents (preferably 16 through 18 years of age).
- c. Bexsero® and Trumenba® are not interchangeable; the same vaccine product must be used for all doses in a series.
- d. Based on available data and expert opinion, Bexsero® and Trumenba® may be administered concomitantly with other vaccines indicated for this age, but at a different anatomic site, if feasible.
- e. Bexsero® and Trumenba® are licensed as both a 2-dose (0.5 mL at 0 and 6 months) series and a 3-dose (0.5 mL at 0, 1-2, and 6 months)

- i. For persons at increased risk for meningococcal disease and for use during serogroup B meningococcal disease outbreaks, the 3-dose schedule **should** be administered to provide earlier protection and maximize short-term immunogenicity. However, if the second dose of MenB vaccine is administered at an interval of ≥ 6 months, a third dose does not need to be administered.
 - ii. For healthy adolescents who are not at increased risk for meningococcal disease, administer 2 doses of MenB vaccine at 0 and 6 months with shared decision making. If the second dose is administered earlier than 6 months after the first dose, a third dose should be administered at least 4 months after the second dose.
 - iii. The 3-dose series (0, 1-2, 6 months) **may** also be used to optimize rapid protection for healthy individuals who initiate the vaccine series less than 6 months prior to increased risk (e.g., within 6 months of college, military recruits).
 - f. MenB vaccines are not recommended for persons who travel to or reside in countries where meningococcal disease is epidemic or hyperendemic because the risk for meningococcal disease in these countries generally is not caused by serogroup B.
 - g. Before administering MenB vaccines, providers should consult the package insert for precautions, warnings, and contraindications.
- (3) MenABCWY (conjugate and recombinant protein) pentavalent vaccine (Penbraya®):
- a. Approved for ages 10-25 years when both MenACWY and MenB vaccines are indicated at the same visit.
 - b. This includes healthy individuals ages 16-23 years who choose to receive MenB vaccination and individuals 10 years and older who are at increased risk of meningococcal disease (persistent

complement deficiencies, complement inhibitor use or anatomic asplenia) and are due for both vaccines.

- c. Contains 2 sterile components: a lyophilized MenACWY component and a MenB component (Trumenba®).
- d. As MenB vaccines are not interchangeable, if a patient receives the MenABCWY vaccine (Penbraya®), then Trumenba® should be used for additional MenB doses when MenACWY is not indicated. They may receive any MenACWY vaccine when MenB is not indicated.
- e. The minimum interval between MenABCWY vaccines (Penbraya®) is 6 months.

3. References.

- a. Mbaeyi SA, Bozio CH, Duffy J, et al. Meningococcal Vaccination: Recommendations of the Advisory Committee on Immunization Practices, United States, 2020. MMWR Recomm Rep 2020;69(No. RR-9):1–41. <http://dx.doi.org/10.15585/mmwr.rr6909a->
- b. Centers for Disease Control and Prevention. Meningococcal Vaccine Recommendations, 2024. Retrieved from <https://www.cdc.gov/vaccines/vpd/mening/hcp/recommendations.html>
- c. Centers for Disease Control and Prevention. Meningococcal Disease; Manual for the Surveillance of Vaccine Preventable Diseases, 2022. Retrieved from <https://www.cdc.gov/vaccines/pubs/surv-manual/chpt08-mening.html>
- d. Multiple resources (e.g., package inserts, Vaccine Information Statements) assembled by DHA-IHD: www.health.mil/meningococcal.

North Atlantic Region Vaccine Safety Hub
Approved: Deputy Chief, Immunization Healthcare Division
(877) 438-8222 (DSN 761-4245)