


Travel and Military Specific Vaccine Labels

Storage and Beyond-Use Date Tracking Labels

Proper storage and handling practices play a very important role in the safety and efficacy of vaccines in protecting individuals and communities from vaccine-preventable diseases.

DHA-IHD recommends providers follow vaccine storage and handling best practices outlined in CDC's Vaccine Storage and Handling Toolkit, including:

- Store each type of vaccine or diluent, if applicable, in its original packaging and in a separate container or bin within the storage unit.
- Use storage labels to help staff easily identify vaccines. The following labels are examples that may be used to help organize vaccines.
- Label shelves and containers to clearly identify where each vaccine and diluent, if applicable, is stored.
- Store vaccines and diluents with similar packaging or names or with pediatric and adult formulations on different shelves.
- If possible, color-code labels (e.g., one color for pediatric and another for adult vaccines).



Adenovirus Type 4 and Type 7

Store between 2°C and 8°C (36°F and 46°F)

Ages: 17 through 50 years

Presentation: Two enteric-coated tablets
((1) Type-4 and (1) Type-7))

Do Not Freeze

Keep bottle tightly closed and protect from moisture

Do NOT remove desiccant from bottle

Biothrax (Anthrax)

Store between 2°C and 8°C (36°F and 46°F)

Ages: 18 through 65 years

Presentation: Multi-dose vial

Do Not Freeze

Beyond Use Time: Once the stopper of the multi-dose vial has been pierced, discard within 28 days

Imovax (Rabies)

Store between 2°C and 8°C (36°F and 46°F)

Ages: All age groups

Presentation: Single-dose vial freeze-dried vaccine and manufacturer-filled syringe diluent

Do Not Freeze

Beyond Use Time: Use immediately after reconstitution

Ixchiq (Chikungunya)

Store between 2°C and 8°C (36°F and 46°F)

Ages: 18 years and older

Presentation: Single-dose vial freeze-dried vaccine and manufacturer-filled syringe diluent

Protect From Light

Do Not Freeze

Ixiaro (Japanese Encephalitis)

Store between 2°C and 8°C (36°F and 46°F)

Ages: 2 months and older

Presentation: Manufacturer-filled syringe

Protect From Light

Do Not Freeze

RabAvert (Rabies)

Store between 2°C and 8°C (36°F and 46°F)

Ages: All age groups

Presentation: Single-dose vial freeze-dried vaccine and manufacturer-filled syringe diluent

Protect From Light

Do Not Freeze

Beyond Use Time: Use immediately after reconstitution

Ticovac (Tick-borne Encephalitis)

Store between 2°C and 8°C (36°F and 46°F)

Ages: 1 year and older

Presentation: Manufacturer-filled syringe

Protect From Light

Do Not Freeze

Typhim Vi (Typhoid)

Store between 2°C and 8°C (36°F and 46°F)

Ages: 2 years and older

Presentation: Manufacturer-filled syringe OR multi-dose vial

Do Not Freeze

Vaxchora (Cholera-Oral)

Store between 2°C and 8°C (36°F and 46°F)

Ages: 2 through 64 years

Presentation: Single-dose carton containing two packets (buffer packet and active component packet)

Do Not Freeze

Protect from light and moisture at all times

Beyond Use Time: After reconstitution, consume within 30 minutes

Vivotif (Typhoid-Oral)

Store between 2°C and 8°C (36°F and 46°F)

Ages: 6 years and older

Presentation: 4 capsules in blister pack

Do Not Freeze

YF-VAX (Yellow Fever)

Store between 2°C and 8°C (36°F and 46°F)

Ages: 9 months and older

Presentation: Single-dose vial lyophilized vaccine and single-dose vial diluent

Do Not Freeze

Beyond Use Time: After reconstitution, administer immediately or store between 2°C and 8°C (36°F and 46°F) and use within 60 minutes

Store between 2°C and 8°C (36°F and 46°F)