TABLE 3. Numbers and Percentages of the Most Frequent Diagnoses During Ambulatory Visits Among Women by ICD-10 Major Diagnostic Category, Active Component, U.S. Armed Forces, 2023

%a
30 14.8 39 10.9 30 3.2 30 3.2 30 3.2 30 8.5 31 7.6 30 7.3 31 6.7 31 5.7
99 10.9 13 10.2 19 3.2 78 3.1 33 70 8.5 61 7.6 97 7.3 13 6.7 91 5.7
13 10.2 39 3.2 78 3.1 103 70 8.5 61 7.6 97 7.3 13 6.7 91 5.7
3.2 3.3 3.1 3.3 3.0 3.0 3.0 3.0 3.0 3.0 3.0
78 3.1 93 70 8.5 97 7.6 97 7.3 13 6.7 91 5.7
70 8.5 61 7.6 97 7.3 13 6.7 91 5.7
70 8.5 61 7.6 67 7.3 13 6.7 61 5.7
51 7.6 97 7.3 13 6.7 91 5.7
97 7.3 13 6.7 91 5.7
3 6.7 91 5.7
91 5.7
0
9 12.3
85 8.2
5.2
1 4.5
3.9
9
16.8
54 7.5
3.4
2.9
18 2.7
)6
8 39.7
4 14.2
50 7.9
92 5.7
3.3
29
11 6.9
18 6.4
39 5.6
90 5.5
11 3.4
37
00 7.9
3 4.5
1 4.2
9 2.6
9 2.6 0 2.3
0 2.3
0 2.3 8
2.3 58 27 9.6
2.3 58 27 9.6 26 8.1
3 5 6 4 4 7 7 5 9 6 2 1 4 3 9 1 3

 $Abbreviations: ICD-10, International\ Classification\ of\ Diseases,\ 10th\ Revision;\ No.,\ number;\ NEC,\ not\ elsewhere\ classified.$

^a Percentage of the total number of hospitalizations within the diagnostic category.

^b Other factors influencing health status and contact with health services (excluding pregnancy-related).

[°]Z-code data are provisional but current as of April 18, 2024.